

Proposed Decision Memo for Transcutaneous Electrical Nerve Stimulation for Chronic Low Back Pain (CAG-00429N)

Decision Summary

CMS proposes coverage for Transcutaneous Electrical Nerve Stimulation (TENS) for chronic low back pain (CLBP) only when all of the following conditions are met.

A. For the purposes of this decision CLBP is defined as:

- a. an episode of low back pain that has persisted for three months or longer; and
- b. is not the result of certain well-defined diseases that may contribute to low back pain but which are not primarily low back syndromes.

For example, there are cancers that, through metastatic spread to the spine or pelvis, may elicit pain in the lower back as a symptom. Certain systemic diseases, e.g. rheumatoid arthritis, multiple sclerosis etc, manifest many debilitating symptoms of which low back pain is not the primary focus. We believe that the appropriate management of these types of diseases is guided by a systematic strategy aimed at the underlying causes. While TENS may infrequently be used adjunctively in managing the symptoms of these diseases, it is clearly not the primary therapeutic approach.

B. The patient is enrolled in a prospective clinical study that addresses one or more aspects of the following questions in a randomized, controlled design using validated and reliable instruments. This can include randomized crossover designs.

1. Does the use of TENS provide a clinically meaningful reduction in pain in Medicare beneficiaries with CLBP?

2. Does the use of TENS provide a clinically meaningful improvement of function in Medicare beneficiaries with CLBP?
3. Does the use of TENS provide a clinically meaningful reduction in other medical treatments or services used in the medical management of CLBP?

The study must adhere to the following standards of scientific integrity and relevance to the Medicare population:

- a. The principal purpose of the clinical study is to test whether TENS potentially improves the participants' health outcomes.
- b. The clinical study is well supported by available scientific and medical information or it is intended to clarify or establish the health outcomes of interventions already in common clinical use.
- c. The clinical study does not unjustifiably duplicate existing studies.
- d. The study design is appropriate to answer the research question being asked in the study.
- e. The clinical study is sponsored by an organization or individual capable of successfully executing the proposed study.
- f. The clinical study is in compliance with all applicable Federal regulations concerning the protection of human subjects found at 45 CFR Part 46.
- g. All aspects of the clinical study are conducted according to appropriate standards of scientific integrity (see <http://www.icmje.org>).
- h. The clinical study has a written protocol that clearly addresses, or incorporates by reference, the standards listed here as Medicare requirements for CED coverage.
 - i. The clinical study is not designed to exclusively test toxicity or disease pathophysiology in healthy individuals.
 - j. The clinical study is registered on the ClinicalTrials.gov website by the principal sponsor/investigator prior to the enrollment of the first study subject.
- k. The clinical study protocol specifies the method and timing of public release of all pre-specified outcomes to be measured including release of outcomes if outcomes are negative or study is terminated early. The results must be made public within 24 months of the end of data collection. If a report is planned to be published in a peer reviewed journal, then that initial release may be an abstract that meets the requirements of the International Committee of Medical Journal Editors

(<http://www.icmje.org>). However a full report of the outcomes must be made public no later than three (3) years after the end of data collection.

- l. The clinical study protocol must explicitly discuss subpopulations affected by the treatment under investigation, particularly traditionally underrepresented groups in clinical studies, how the inclusion and exclusion criteria affect enrollment of these populations, and a plan for the retention and reporting of said populations on the trial. If the inclusion and exclusion criteria are expected to have a negative effect on the recruitment or retention of underrepresented populations, the protocol must discuss why these criteria are necessary.
- m. The clinical study protocol explicitly discusses how the results are or are not expected to be generalizable to the Medicare population to infer whether Medicare patients may benefit from the intervention. Separate discussions in the protocol may be necessary for populations eligible for Medicare due to age, disability or Medicaid eligibility.

Consistent with section 1142 of the Social Security Act, AHRQ supports clinical research studies that CMS determines meet the above-listed standards and address the above-listed research questions.

We are requesting public comments to this proposed decision pursuant to section 1862(l) of the Social Security Act (the Act). After consideration of the public comments and any additional evidence, we will issue a final determination responding to the public comments consistent with §1862(l)(3) of the Act.

[Back to Top](#)

Proposed Decision Memo

CMS proposes coverage for Transcutaneous Electrical Nerve Stimulation (TENS) for chronic low back pain (CLBP) only when all of the following conditions are met.

A. For the purposes of this decision CLBP is defined as:

- a. an episode of low back pain that has persisted for three months or longer; and

- b. is not the result of certain well-defined diseases that may contribute to low back pain but which are not primarily low back syndromes.

For example, there are cancers that, through metastatic spread to the spine or pelvis, may elicit pain in the lower back as a symptom. Certain systemic diseases, e.g. rheumatoid arthritis, multiple sclerosis etc, manifest many debilitating symptoms of which low back pain is not the primary focus. We believe that the appropriate management of these types of diseases is guided by a systematic strategy aimed at the underlying causes. While TENS may infrequently be used adjunctively in managing the symptoms of these diseases, it is clearly not the primary therapeutic approach.

B. The patient is enrolled in a prospective clinical study that addresses one or more aspects of the following questions in a randomized, controlled design using validated and reliable instruments. This can include randomized crossover designs.

1. Does the use of TENS provide a clinically meaningful reduction in pain in Medicare beneficiaries with CLBP?
2. Does the use of TENS provide a clinically meaningful improvement of function in Medicare beneficiaries with CLBP?
3. Does the use of TENS provide a clinically meaningful reduction in other medical treatments or services used in the medical management of CLBP?

The study must adhere to the following standards of scientific integrity and relevance to the Medicare population:

- a. The principal purpose of the clinical study is to test whether TENS potentially improves the participants' health outcomes.
- b. The clinical study is well supported by available scientific and medical information or it is intended to clarify or establish the health outcomes of interventions already in common clinical use.
- c. The clinical study does not unjustifiably duplicate existing studies.
- d. The study design is appropriate to answer the research question being asked in the study.
- e. The clinical study is sponsored by an organization or individual capable of successfully executing the proposed study.

- f. The clinical study is in compliance with all applicable Federal regulations concerning the protection of human subjects found at 45 CFR Part 46.
- g. All aspects of the clinical study are conducted according to appropriate standards of scientific integrity (see <http://www.icmje.org>).
- h. The clinical study has a written protocol that clearly addresses, or incorporates by reference, the standards listed here as Medicare requirements for CED coverage.
- i. The clinical study is not designed to exclusively test toxicity or disease pathophysiology in healthy individuals.
- j. The clinical study is registered on the ClinicalTrials.gov website by the principal sponsor/investigator prior to the enrollment of the first study subject.
- k. The clinical study protocol specifies the method and timing of public release of all pre-specified outcomes to be measured including release of outcomes if outcomes are negative or study is terminated early. The results must be made public within 24 months of the end of data collection. If a report is planned to be published in a peer reviewed journal, then that initial release may be an abstract that meets the requirements of the International Committee of Medical Journal Editors

(<http://www.icmje.org>). However a full report of the outcomes must be made public no later than three (3) years after the end of data collection.

- l. The clinical study protocol must explicitly discuss subpopulations affected by the treatment under investigation, particularly traditionally underrepresented groups in clinical studies, how the inclusion and exclusion criteria affect enrollment of these populations, and a plan for the retention and reporting of said populations on the trial. If the inclusion and exclusion criteria are expected to have a negative effect on the recruitment or retention of underrepresented populations, the protocol must discuss why these criteria are necessary.
- m. The clinical study protocol explicitly discusses how the results are or are not expected to be generalizable to the Medicare population to infer whether Medicare patients may benefit from the intervention. Separate discussions in the protocol may be necessary for populations eligible for Medicare due to age, disability or Medicaid eligibility.

Consistent with section 1142 of the Social Security Act, AHRQ supports clinical research studies that CMS determines meet the above-listed standards and address the above-listed research questions.

We are requesting public comments to this proposed decision pursuant to section 1862(l) of the Social Security Act (the Act). After consideration of the public comments and any additional evidence, we will issue a final determination responding to the public comments consistent with §1862(l)(3) of the Act.

Bibliography

Akl E, Briel M, You J, Lamontagne F, Gangj A, Cukierman-Yaffe T, Alshurafa M, Sun X, Nerenberg K, Johnston B, Vera C, Mills E, Bassler D, Salazar A, Bhatnagar N, Busse J, Khalid Z, Walter S, Cook D, Schunemann H, Altman D, Guyatt g. LOST to follow-up Information in Trials (LOST – IT): a protocol on the potential impact. *Trials*. 2009, 10:40.

American Medical Directors Association (AMDA). Pain management in the long-term care setting. Columbia (MD): American Medical Directors Association (AMDA). 2009:48.

American Pain Society. Guidelines for the Use of Chronic Opioid Therapy in Chronic Noncancer Pain: Evidence Review. Glenview, IL: American Pain Society. 2009.

Barker K, Elliot C, Sackley C, Fairbank J. Treatment of chronic back pain by sensory discrimination training. A Phase I RCT of a novel device (FairMed) vs. TENS. *BCM Musculoskeletal Disorders*. 2008;9:97.

Bennett M, Hughes N., and Johnson M. Methodological quality in randomised controlled trials of transcutaneous electric nerve stimulation for pain: Low fidelity may explain negative findings. *Pain*, 2011;152:1226-1232.

Bjordal J. Time for a paradigm shift in pain treatment: Reassessing transcutaneous electrical nerve stimulation (TENS). *Pain*. 2011; 152: 1213-1214.

Bjordal J, Johnson M, Ljunggreen A. Transcutaneous electrical nerve stimulation (TENS) can reduce postoperative analgesic consumption. A meta-analysis with assessment of optimal treatment parameters for postoperative pain. *European Journal of Pain*. 2003; 7:181-188.

Bloodworth D, Calvillo O, Smith K, and Grabois M. Chronic Pain Syndromes: Evaluation and Treatment in *Physical Medicine and Rehabilitation* [ed:RL Braddom]. W.B. Saunders, 2000: 913-933.

Boutron I, Moher D, Altman D, Schulz K, and Philippe R. Methods and Processes of the CONSORT Group: Example of an Extension for Trials Assessing Nonpharmacologic Treatments. *Annals of Internal Medicine*, 2008;148(4):w60-w66.

Brosseau L, Milne S, Robinson V, Marchand S, Shea B, Wells G, Tugwell P. Efficacy of the Transcutaneous Electrical Nerve Stimulation for the Treatment of Chronic Low Back Pain. A Meta-Analysis. *Spine*. 2002;27(6):596-603.

Chabel C, Fishbain D, Weaver M, Wipperman Heine, L. Long-Term transcutaneous electrical nerve stimulation (TENS) use: impact on medication utilization and physical therapy costs. *Clinical Journal of Pain*. 1998; 14:66-73.

Cheing G, Hui-Chan W. Y. Transcutaneous Electrical Nerve Stimulation: Nonparallel Antinociceptive Effects on Chronic Clinical pain and Acute Experimental Pain. *Archives of Physical Medicine and Rehabilitation*. 1999;80:305-312.

Chou R, Fanciullo G, Fine P, Adler J, Ballantyne J, Davies P, Donovan M, Fishbain D, Foley K, Fudin J, Gilson A, Kelter A, Mauskop A, O'Connor P, Passik S, Pasternak G, Portenoy R, Rich B, Roberts R, Todd K, Miaskowski C. American Pain Society-American Academy of Pain Medicine Opioids Guidelines Panel. Clinical guidelines for the use of chronic opioids therapy in chronic noncancer pain. *Journal of Pain*. 2009;10(2): 113-130.

Chou R, Huffman L. Nonpharmacologic Therapies for Acute and Chronic Low Back pain: A Review of the Evidence for an American Pain Society/American College of Physicians Clinical Practice Guideline. *Annals of Internal Medicine*. 2007;147:492-504.

Chou R, Shekelle P. Will this Patient Develop Persistent Disabling Low Back Pain? *Journal of the American Medical Association*. 2010; 303(13):1295-1302.

Cochrane Collaboration open learning material: Further issues in meta-analysis, Intention to treat issues. Retrieved January 11, 2012 from <http://www.cochrane-net.org/openlearning/HTML/mod14-4.htm>.

Davidhizar R., Giger J. A review of the literature on care of clients in pain who are culturally diverse. *International Nursing Review*. 2004;51:47-55.

DeSantana J, Sluka K. Central mechanisms in the maintenance of chronic widespread noninflammatory muscle pain. *Current Pain and Headache Reports*. 2008;12:338-343.

DeSantana J, Walsh D, Vance C, Rakel B, Sluka K. Effectiveness of transcutaneous electrical nerve stimulation for treatment of hyperalgesia and pain. *Current Rheumatology Reports*. 2008;10:492-499.

Deyo R, Walsh N, Martin D, Schoenfeld L, Ramamurthy S. A Controlled Trial of Transcutaneous Electrical Nerve Stimulation (TENS) and Exercise for Chronic Low Back Pain. *New England Journal of Medicine*. 1990;322(23):1627-1634.

Deyo R, Walsh N, Martin D, Schoenfeld L, Ramamurthy S. Can Trials of Physical Treatments Be Blinded? The Example of Transcutaneous Electrical Nerve Stimulation for Chronic Pain. *American Journal of Physical Medicine & Rehabilitation*. 1990;69(1):6-10.

Dubinsky R, Miyasaki J. Assessment: Efficacy of transcutaneous electric nerve stimulation in the treatment of pain in neurologic disorders (an evidence-based review). Report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology*. 2010;74:173-176.

Elkiss M, Ingall J, Dooley J. Pilot study on the effects of transcutaneous electrical nerve stimulation on the level of plasma beta-endorphin. *Journal of the American Osteopathic Association*. 1984;84(suppl. 1):144-147.

Fishbain D, Chabal C, Abbott A. , Heine L, Cutler R. Transcutaneous Electrical Nerve Stimulation (TENS). *Clinical Journal of Pain*. 1996;12(3):201-14.

Flowerdew M, Gadsby J. A review of the treatment of chronic low back pain with acupuncture-like transcutaneous electrical nerve stimulation and transcutaneous electrical nerve stimulation. *Complementary Therapies in Medicine*. 1995;5:193-201.

Gadsby J, Flowerdew M. Transcutaneous electrical nerve stimulation and acupuncture-like transcutaneous electrical nerve stimulation for chronic low back pain (withdrawn). The Cochrane Database of Systematic Reviews 1997, Issue 1. Art. No. :CD000210. DOI:10.1002/14651858. CD000210.

Grady D, Berkowitz S, Katz M. Opioids for Chronic Pain. *Archives of internal Medicine*. 2011;171(16).

Grant D, Bishop-Miller J, Winchester D, Anderson M, Faulkner S. A randomized comparative trial of acupuncture versus transcutaneous electrical nerve stimulation for chronic back pain in the elderly. *Pain*. 1999; 82: 9-13.

Grazio S, Markuiincic B, Muraja S, Grubisic F, Nemcic T, Matijevic V, Skala H, Kasun B, Susak V, Koprivnak V, Trgovec I. Efficacy of low-level laser therapy and transcutaneous electrical nerve stimulation (TENS) therapy on level of pain, global assessment of the disease and function in chronic low-back pain. *Clinical and Experimental Rheumatology*. 2009;27(5):725-726.

Griffin J, Kraft G. Patients with chronic pain treatment by transcutaneous electrical nerve stimulation: Comparison of F = 250 Hz or less versus F = 1,000 Hz. *International Journal of Rehabilitation Research*. 1981;4(2):230-231.

Harris G, Susman J. Managing musculoskeletal complaints with rehabilitation therapy: Summary of the Philadelphia Panel evidence-based clinical practice guidelines on musculoskeletal rehabilitation interventions. *The Journal of Family Practice*. 2002;51(12):1042-1046.

Itoh K, Itoh S, Katsumi Y, Kitakoji H. A pilot study on using acupuncture and transcutaneous electrical nerve stimulation to treat chronic non-specific low back pain. *Complementary Therapies in Clinical Practice*. 2009;15:22-25.

Jarzem P, Harvey E, Arcaro N, Kaczorowski J. Transcutaneous electrical nerve stimulation (TENS) for chronic low back pain. *Journal of Musculoskeletal Pain*. 2005;13(2):3-9.

Jarzem P, Harvey E, Arcaro N, Kaczorowski J. Transcutaneous electrical nerve stimulation (TENS) for short-term treatment of low back pain – Randomized double blind crossover study of sham versus conventional TENS. *Journal of Musculoskeletal Pain*. 2005;13(2):11-17.

Johnsen A, Sonne N. Transcutaneous Electrical Nerve Stimulation (tens) shows pain relief in patients with chronic pain. *European Journal of Pain*. 2009;13(suppl1):S252

Johnson M., Din A. Ethnocultural differences in the analgesic effects of placebo transcutaneous electrical nerve stimulation on cold-induced pain in healthy subjects: a preliminary study. *Complementary Therapies in Medicine*. 1997;5:74-79.

Johnson M, Ashton C, Thompson J. Long term use of transcutaneous electrical nerve stimulation at New Castle Pain Relief Clinic. *Journal of the Royal Society of Medicine*, 1992;85:267-268.

Johnson M, Ashton C, Thompson J. An in-depth study of long-term users of transcutaneous electrical nerve stimulation (TENS). Implications for clinical use of TENS. *Pain*. 1991;44:221-229.

Johnson M, Martinson M. Efficacy of electrical nerve stimulation for chronic musculoskeletal pain: A meta-analysis of randomized controlled trials. *Pain*. 2007;130:157-165.

Karla A, rban M, Sluka K. Blockade of opioid receptors in rostral ventral medulla prevents antihyperalgesia produced by transcutaneous electrical nerve stimulation (TENS). *Journal of Pharmacology and Experimental Therapeutics*. 2001;298: 257-263.

Keller A, Hayden J, Bombardier C, van Tulder M. Effect sizes of non-surgical treatments of non-specific low-back pain. *European Spine Journal*. 2007;16:1776-1788.

Kerns R, Kassirer M, Otis J. Pain in multiple sclerosis: a biopsychosocial perspective. *Journal of Rehabilitation Research and Development*. 2002;39:225-232.

Kerr D, Walsh D, Baxter D. Acupuncture in the Management of Chronic Low Back Pain: A Blinded Randomized Controlled Trial. *The Clinical Journal of Pain*. 2003;19(6):364-370.

Khadilkar A. Transcutaneous Electrical Nerve Stimulation for the Treatment of Chronic Low Back Pain: A Systematic Review. *Spine*. 2005;30(23):2657-2666.

Knight K, Draper, D. Therapeutic Modalities: The Art and Science with Clinical Activities Manual. Lippincott Williams & Wilkins. 2007.

Kofotolis N, Vlachopoulos S, Kellis E. Sequentially allocated clinical trial of rhythmic stabilization exercises and TENS in women with chronic low back pain. *Clinical Rehabilitation*. 2008;22:99-111.

Koldas D, Sonel T, Kurtais Y, Atay M. Comparison of three different approaches in the treatment of chronic low back pain. *Clinical Rheumatology*. 2008;7:873-81.

Lehmann T, Russell D, Spratt K. The impact of patients with nonorganic physical findings on a controlled trial of transcutaneous electrical nerve stimulation and electoracupuncture. *Spine*. 1983;8(6):625-634.

Lehmann T, Russell D, Spratt K, Colby H, Liu Y, Fairchild M, Christensen S. Efficacy of Electroacupuncture and TENS in the Rehabilitation of Chronic Low Back Pain Patients. *Pain*. 1986;26:277-290.

Léonard, G, Cloutier C, Marchand S. Reduced analgesic effect of acupuncture-like tens but not conventional tens in opioid-treated patients. *Journal of Pain*. 2011;12(2):213-221.

Liebano R, Rakel B, Vance C, Walsh D, Sluka K. An investigation of the development of analgesic tolerance to TENS in humans. *Pain*, 2011;152(2):335-342.

Lin, M, Lin, M, Fen J, Lin W, Lin C, Chen P. A comparison between pulsed radiofrequency and electro-acupuncture for relieving pain in patients with chronic low back pain. *Acupuncture & electro-therapeutics research*. 2010; 35(3-4):133-46.

Machado L, Kamper S, Herbert R, Maher C, McAuley J. Analgesic effects of treatments for non-specific low back pain: a meta-analysis of placebo-controlled randomized trials. *Rheumatology*. 2009;46:520-527.

Mannheimer J, Lampe G. Specific stimulation modes: A literature review. In Mannheimer JS, Lampe GN (eds): Clinical Transcutaneous Electrical Nerve Stimulation. 1984. FA Davis Company, Philadelphia, PA, 338-348.

Marchand S, Charest J, Li J, Chenard J, Lavignolle B, Laurencelle L. Is TENS purely a placebo effect? A controlled study on chronic low back pain. *Pain*. 1993;54:99-106.

Martinson, M, Johnson M. Cost-effectiveness of Treatments for Low Back Pain. *Practical Pain Management* 2011; 23:32-37.

McIntosh G, Hall H. Low back pain (chronic). *Clinical Evidence*. 2008;10:1116.

Melzak R, Vetere P, Finch L. Transcutaneous Electrical Nerve Stimulation for Low Back Pain. A Comparison of TENS and Massage for Pain and Range of Motion. *Physical Therapy*. 1983;63(4);489-493.

Milne S, Welch V, Brosseau L, Saginur M, Shea B, Tugwell P, Wells G. Transcutaneous electrical nerve stimulation (TENS) for chronic low-back pain. *The Cochrane Database of Systematic Reviews*, Issue. Art. No. :CD003008. DOI: 10. 1002/14651858. CD003008.

Minkoff J, Harris G, Susman J. Managing musculoskeletal complaints with rehabilitation therapy: Summary of the Philadelphia Panel evidence-based clinical practice guidelines on musculoskeletal rehabilitation interventions. *Journal of Family Practice*. 2002;51(12):1042-1046.

Moran F, Leonard T, Hawthorne S, Hughes C, Crum-Gardner E, Johnson M, Rakel B, Sluka K, Walsh D. Hypoalgesia in Response to Transcutaneous Electrical Nerve Stimulation (TENS) Depends on Stimulation Intensity. *Journal of Pain*. 2011; 12: 929-935.

Negrini S, Zaina F, Romano M, Atanasio S, Fusco C, Trevisan C. “Rehabilitation of Lumbar Spine Disorders: An Evidence – Based Clinical Practice Approach” in DeLisa’s Physical Medicine and Rehabilitation, Principles and Practice. Frontera et. al. (eds.) Wolters Kluwer/Lippincott Williams and Wilkins. 2010:837 – 882.

Noble J, Henderson G, Cramp A, Walsh D and Lowe A. The effect of interferential therapy upon cutaneous blood flow in humans. *Clinical Physiology*. 2008;20:2-7.

Novak S, Nemeth W. How clinically relevant is a meta-analysis of electrical nerve stimulation when based on heterogeneous disease states? *Pain*. 2007;131(1-2):228-229.

Ostelo RWJG, de Vet HCW. Clinically important outcomes in low back pain. *Best Practice and Research Clinical Rheumatology*. 2005; 19(4): 593-607.

Poitras S, Brosseau L. Evidence-informed management of chronic low back pain with transcutaneous electrical nerve stimulation, interferential current, electrical muscle stimulation, ultrasound, and thermotherapy. *The Spine Journal*. 2008;8:226-233.

Rakel B, Cooper N, Adams H, Messer B, Frey Law L, Dannen D, Miller C, Polehna A, Ruggle R, Vance C, Walsh D, Sluka K. A New Transient Sham TENS Device Allows for Investigator Blinding While Delivering a True Placebo Treatment. *Journal of Pain*. 2010;11: 230-238.

Rakel B, Frantz R. Effectiveness of transcutaneous electrical nerve stimulation on postoperative pain with movement. *Journal of Pain*. 2003;4:455-464.

Rathmell JP. A 50-Year Old Man with Chronic Low Back Pain. *Journal of the American Medical Association*. 2008; 299(17):2066-2077.

Reigelman R. *Studying a Study & Testing a Test*. How to read the medical evidence. (5th edition) Lippincott Williams and Wilkins. 2005.

Rushton, D. Electrical stimulation in the treatment of pain<. *Disability and Rehabilitation*. 2002;24(8):407-15.

Seco J, Kovacs F, Urrutia G. The efficacy, safety, effectiveness, and cost-effectiveness of ultrasound and shock wave therapies for low back pain: a systematic review. *The Spine Journal*. 2011:1-12.

Shimoji K, Takahashi N, Nishio Y, Koyanagi M, Aida S. Pain Relief by Transcutaneous Electrical Nerve Stimulation With Bidirectional Modulated Sine Waves in Patients With Chronic Back Pain: A Randomized, Double-Blind, Sham-Controlled Study. *Neuromodulation: Technology at the Neural Interface*. 2007;10(1):42-51.

Sluka K (ed). Mechanisms and Management of Pain for the Physical Therapist. IASP Press. 2009.

Stanos S, Tyburski M, Harden R. Management of Chronic Pain in *Physical Medicine and Rehabilitation* [ed:RL Braddom]. Saunders Elsevier, 2007: 951-988.

Thorsteinsson G, Stonnington H, Stillwell G, Elveback L. Transcutaneous Electrical Stimulation: A Double-Blind Trial of its Efficacy for Pain. *Archives of Physical Medicine & Rehabilitation*. 1977;58.

Topuz O, Ozfidan E, Ozgen M, Ardic F. Efficacy of transcutaneous electrical nerve stimulation and percutaneous neuromodulation therapy in chronic low back pain. *Journal of Back and Musculoskeletal Rehabilitation*. 2004;17(3-4):127-133.

Tsukayama H, Yamashita H, Amagai H, Tanno Y. Randomised Controlled Trial Comparing the Effectiveness of Electroacupuncture and TENS for Low Back Pain: A Preliminary Study for a Pragmatic Trial. *Acupuncture in Medicine*. 2002;20(4):175-180.

Tulgar M, McGlone F, Bowsher D, Miles J. Comparative effectiveness of different stimulation modes in relieving pain. Part I. A pilot study. *Pain*. 1991(a);47:151-155.

Tulgar M, McGlone F, Bowser D, Miles J. Comparative effectiveness of different stimulation modes in relieving pain. Part II. A double-blind controlled long-term clinical trial. *Pain*. 1991(b);47:157-162.

van Middelkoop M, Rubinstein S, Kuijpers T, Verhagen A, Ostelo R, Koes B, van Tulder M. A systematic review on the effectiveness of physical and rehabilitation interventions for chronic non-specific low back pain. *European Spine Journal*. 2011;20:19-39.

van Tulder M, Gherkin D, Berman B, Lao L, Koes B. The Effectiveness of Acupuncture in the Management of Acute and Chronic Low Back Pain: A Systematic Review Within the Framework of the Cochrane Collaboration Back Review Group. *Spine*. 1999;24(11):1113-1123.

van Tulder M, Gherkin D, Berman B, Lao L, Koes B. Acupuncture for low-back pain. The *Cochrane Database of Systematic Reviews*, Issue. Art. No. :CD001351. DOI: 10.1001/14651858. CD001351.

Walsh N, Santa Maria D, Eckmann M. Treatment of the Patient with Chronic Pain in Physical Medicine and Rehabilitation Principles and Practice [Frontera WR, editor in chief]. Wolters Kluwer/Lippincott Williams and Wilkins. 2010.

Warke K, Al-Smadi J, Baxter D, Walsh D, Lowe-Strong A. Efficacy of Transcutaneous Electrical Nerve Stimulation (TENS) for Chronic Low-back Pain in a Multiple Sclerosis Population. A Randomized, Placebo-controlled Clinical Trial. *Clinical Journal Pain*. 2006;22(9):812-819.

Weinstein SM, Herring SA, Standaert CJ. Low Back Pain in Physical Medicine and Rehabilitation Principles and Practice [DeLisa JA, editor in chief]. Lippincott Williams and Wilkens. 2005;653-678.

Winnem M, Staff P. Transcutaneous electrical stimulation in the treatment of pain. *Tidsskrift for den Norske Laegeforening*. 1982;102(29):1511-1513+1520.

Yeung, C, Leung, M, Chow D. The use of electro-acupuncture in conjunction with exercise for the treatment of chronic low-back pain. *Journal of Alternative & Complementary Medicine*. 2003;9(4):479-90.

Yip Y, Sonny T, Wu K. An experimental study comparing the effects of combined transcutaneous acupoint electrical stimulation and electromagnetic millimeter waves for spinal pain Hong Kong. *Complementary Therapies in Clinical Practice*. 2007;13(1):4-14.

Yokoyama M, Sun X, Oku S, Taga N, Sato K, Mizobuchi S, Takahashi T, Morita K. Comparison of Percutaneous Electrical Nerve Stimulation with Transcutaneous Electrical Nerve Stimulation for Long-Term Pain Relief in Patients with Chronic Low Back Pain. *Anesthesia & Analgesia*. 2004;98:1552-6.

Zambito A, Bianchini D, Gatti D, Rossini M, Adami S, Viapiana O. Inferential and horizontal therapies in chronic low back pain due to multiple vertebral fractures: a randomized, double blind, clinical study. *Osteoporosis International*. 2007;18(11):1541-5.

Zambito A, Bianchini D, Gatti D, Viapiana O, Rossini M, Adami S. Inferential and horizontal therapies in chronic low back pain: a randomized, double blind, clinical study. *Clinical and Experimental Rheumatology*. 2006;24(5):534-9.

[Back to Top](#)